Journal of Immuological Methods, 121 (1989) 298
Elsevier

Contents to volume 120 (1989)

Number 1 – 6 July 1989

Research reports

Enhanced fluorescence in indirect immunophenotyping by the use of fluorescent liposomes
A.G. Gray, J. Morgan, D.C. Linch and E.R. Huehns (U.K.) 1

A novel and inexpensive source of allophycocyanin for multicolor flow cytometry
T.M. Jung and M.O. Dailey (U.S.A.) 9

Assessment of human allospecific IL-2-secreting helper T lymphocytes in limiting dilution cultures using restimulation and split well analysis
M. Theobald, T. Hoffmann and W. Heit (F.R.G.) 19

Recovery of functional human lymphocytes from Leukotrap filters
R.E. Longley and D. Stewart (U.S.A.) 33

Analysis of cytotoxic T cell precursor frequencies directed against individual HLA-A and -B alloantigens
L. Zhang, S.G. Li, B. Vandekerckhove, A. Termijtelen, J.J. Van Rood and F.H.J. Claas (The Netherlands) 39

Immunodetection of human tumour-associated cell-bound antigens by human monoclonal antibodies
J.R.D. Tuson, D.A. Jacob, E.W. Pascoe and S.J.Y. Saxby (U.K.) 47

Production of multilamellar, small unilamellar and reverse-phase liposomes containing house dust mite allergens. Potential adjuvants in the immunotherapy of allergic disease
A.S. McWilliam and G.A. Stewart (Australia) 53

Biotinylation: a simple method for labelling complement component C8 with preservation of functional activity
S. Bhakdi, M. Roth and F. Hugo (F.R.G.) 61

In vitro detection of lymphokines produced by in vivo activated lymphocytes
K.M. Mohler and L.D. Butler (U.S.A.) 67

Evaluation of methods for isolating human peripheral blood monocytes. Studies on chemotactic locomotion and other functional characteristics
L.N. Islam and P.C. Wilkinson (U.K.) 75

Interference of complement with the binding of carcinoembryonic antigen to solid-phase monoclonal antibodies
O.P. Börmer (Norway) 85

Determination of human antibodies to specific antigens of cytomegalovirus using an antigen capture immunoassay
L.H.M. Van der Voort, J. Van Zanten, L.F.M.H. De Leij and T.H. The (The Netherlands) 95

Rapid method for the isolation of neutrophils in high yield without the use of dextran or density gradient polymers
P. Eggleton, R. Gargan and D. Fisher (U.K.) 105

Natural killer function in flow cytometry. II. Evaluation of NK lytic activity by means of target cell morphological changes detected by right angle light scatter
M. Vitale, L.M. Neri, S. Comani, E. Falcieri, R. Rizzoli, R. Rana and S. Papa (Italy) 115

Homogeneous immunoassay for α_2 plasmin inhibitor (α_2PI) and α_2PI-plasmin complex. Application of a sandwich liposome immune lysis assay (LILA) technique
K. Hosoda and T. Yasuda (Japan) 121

The effects of aromatic and aliphatic maleimide crosslinkers on anti-CD5 ricin immunotoxins
D.E. Myers, F.M. Uckun, S.E. Swaim and D.A. Vallera (U.S.A.) 129

Letters to the editors

Comments on 'Measurement of tumour reactive antibody and antibody conjugate by competition, quantitated by flow cytometry', by Robins et al. (1986) J. Immunol. Methods 90, 165–172
P.A. Underwood (Australia) 143

Reply to the letter of P.A. Underwood
R.A. Robins and R.R. Laxton (U.K.) 145

Book reviews 147

Announcement 149

Notes to authors 150

0022-1759/89/$03.50 © 1989 Elsevier Science Publishers B.V. (Biomedical Division)

U·M·I

INCORRECT VOLUME NUMBER, SHOULD
READ VOLUME 121.

Number 2 – 26 July 1989

Research reports

An enzyme immunoassay for the determination of low levels of IgA in human serum
M. Haun and S. Wasi (Canada) 151

Influence of adjuvants on the quantity, affinity, isotype and epitope specificity of murine antibodies
J.S. Kenney, B.W. Hughes, M.P. Masada and A.C. Allison (U.S.A.) 157

Application of flow cytometric measurement of surface IgG in kinetic analysis of monoclonal antibody synthesis and secretion by murine hybridoma cells
E. Meilhoc, K.D. Wittrup and J.E. Bailey (U.S.A.) 167

Determination of cell-bound Fc receptors using a cell adherence test (CAT)
H. Lemke, G. Huss, J. Bonmann and H. Hansen (F.R.G.) 175

Isolation of encephalitogenic $CD4^+$ T cell clones in the rat. Cloning methodology and interferon-γ secretion
J.D. Sedgwick, I.A.M. MacPhee and M. Puklavec (U.K.) 185

A flow cytometric method for the detection of the development of antibody to Orthoclone OKT3
V.L. Lim, M. Gumbert and M.R. Garovoy (U.S.A.) 197

Staphylococcus aureus phagocytosis. A new cytofluorometric method using FITC and paraformaldehyde
B. Cantinieaux, C. Hariga, P. Courtoy, J. Hupin and P. Fondu (Belgium) 203

Mouse IgG2b monoclonal antibody fragmentation. Preparation and purification of Fab, Fc and Fab/c fragments
S. Demignot, M.C. Garnett and R.W. Baldwin (U.K.) 209

A simple method for the biochemical purification of Ro/SS-A antigen
D.-H. Wu, A. Tavoni, C. Garzelli, R. Neri, C. Vitali and S. Bombardieri (Italy) 219

Requirements for screening recombinant DNA libraries for T cell epitope expression
M. Guttinger, U. Certa, J.R. Pink and F. Sinigaglia (Switzerland) 225

Binding of mitogenic plant lectins to human lymphocytes. Flow cytometric analysis
S. Serke, A. Neubauer and A. Van Lessen (F.R.G.) 231

Generation of a sheep × mouse heterohybridoma cell line (1C6.3a6T.1D7) and evaluation of its use in the production of ovine monoclonal antibodies
J.N. Flynn, G.D. Harkiss and J. Hopkins (U.K.) 237

An enzyme-linked immunosorbent assay (ELISA) for urinary growth hormone suitable for use in the routine laboratory. The Wessex Growth Study
H. Bullen and T.J. Wilkin (U.K.) 247

Optimal conditions for the generation of monoclonal antibodies using primary immunisation of mouse splenocytes in vitro under serum-free conditions
M. De Boer, F.A. Ossendorp, G. Van Duijn, G.H.J. Ten Voorde and J.M. Tager (The Netherlands) 253

Quantitation of the anaphylatoxin C3a in the presence of C3 by a novel sandwich ELISA using monoclonal antibody to a C3a neoepitope
G. Zilow, W. Naser, R. Rutz and R. Burger (F.R.G.) 261

Differential oligoclonal band patterns on polyvinyldifluoride membranes
R.W. Luxton and E.J. Thompson (U.K.) 269

A method for the detection of IgE binding sequences of allergens based on a modification of epitope mapping
B.J. Walsh and M.E.H. Howden (Australia) 275

A fluorescence sandwich ELISA for detecting soluble and cell-associated human interleukin-2
K. Kitamura, K. Matsuda, M. Ide, T. Tokunaga and M. Honda (Japan) 281

Technical note

Non-specific cell binding characteristics of para-magnetic polystyrene microspheres used for antibody-mediated cell selection
W.E. Janssen and A.M. Rios (U.S.A.) 289

Letter to the editors

Enhancement of immunoblot staining using a mixed chromogenic substrate
P.R. Young (U.K.) 295

Announcement 297

Contents to volume 121 (1989) 298

Notes to authors 300